

# ORGANOMETALLIC COMPOUNDS: ITS THERAPEUTIC POTENTIALS TOWARDS CANCER TREATMENT

**Author:** Taiwo Esther Omolola

## ABSTRACT

Organometallic compounds are renowned for their remarkable applications in the field of catalysis, but much less is known about their potential in chemical biology. Indeed, such compounds have long been considered to be either unstable under physiological conditions or cytotoxic. As a consequence, little attention has been paid to their possible utilization for biological purposes. Because of their outstanding physicochemical properties, which include chemical stability, structural diversity and unique photo- and electrochemical properties, however, organometallic compounds can play a leading role in the field of chemical biology. Indeed, remarkable examples of the use of such compounds—namely as enzyme inhibitors and as luminescent agents—have recently been reported. Here, recent advances in the use of organometallic compounds for chemical biology purposes, an area that we define as 'organometallic chemical biology'. We also demonstrate that these recent discoveries are only a beginning and that many other organometallic complexes are likely to be found useful in this field of research soon. Inorganic and organometallic compounds have been used as pesticides. At present, in most developed countries, only copper compounds are used in a significant amount as fungicides. The others have been banned for either toxicological or environmental reasons. It is believed that organotin compounds are still used in some developing countries. Copper compounds if used as directed do not pose health risks, however; the estimated long-term intake in the European Union is not much below the acceptable daily intake when other non-pesticidal sources of copper are considered. Organotin compounds have no negligible acute toxicity. Triphenyltin has immunotoxic and, possibly, carcinogenic effects and cyhexatin/azocyclotin have embryotoxic properties. However, estimated intakes are below reference values.

**Keywords:** organometallic compounds, clinical and industrial applications, Cancer Therapy.

## 1.0 INTRODUCTION

Organometallic compounds are typically discussed in terms of the metals as either main-group; group 1 and 2 that are in the S-block and as well as those P-block which chemical are the heavier elements i.e. the group 13-15 compounds or transition metal compounds, most especially those elements in the D- and F-block, group 3-12.

Knowledge of the structure and concentration of organometallic compounds throughout the ecosystem is important in working out the pathways and mechanisms by which metals distribute themselves throughout the environment. Reactive compounds which are kept in solution in an organic solvent; they form electron rich or anionic carbon atom during their reactions i.e. functions as bases or nucleophiles having a general formula of  $R^-M^+$

Organometallic compounds are reserved for compounds that contain at least one metal carbon bond including alkaline, alkaline earth and transition metals and are sometimes broadened to include metalloids like boron, silicon and selenium. These compounds offer a versatile platform for the design of therapeutics from antimicrobials to anticancer agents.

Alkoxides such as  $C_3H_7O_4Ti$  is not considered organometallic compounds because here, the organic group is bonded to Ti through oxygen and not through carbon; whereas the compound  $C_6H_5Ti(OC_3H_7)_3$  is an organometallic compound because in this compound Ti is bonded to the carbon of the organic group.

Many organometallic compounds are highly toxic, especially those that are volatile, stable, some are spontaneously flammable e.g. lithium, sodium, and aluminum. Most are solids and some are liquids and gases. Its properties depend in large measure on the type of carbon metal bond involved in covalent bonds, multicentre or ionic bonds as well as polarization.

It can be classified also based on the nature of metal carbon bonds as ionic compounds/electropositive metals, covalent organometallic compounds, organometallic compounds involving p-bonds, carbene/carbyne compounds and transition metal organometallic compounds.

Examples of organometallic compounds are Zeise salt prepared by the Danish pharmacist William c. Zeise in 1827 has no way of knowing the structure of his novel compound, but it is now known that the ethylene molecules are connected to the center platinum (Pt) atom through both carbon atoms. Three chlorine (Cl) atoms are bonded to the Platinum atom. The potassium

ion  $K^+$  present to balance charge  $K(PtCl_3(C_2H_4))$ . Other examples, ferrocene  $(C_2H_5)_2$  and tetraethyl lead  $Pb(C_2H_5)_4$ .

The significance of organometallic intensities rests in the way that they have a vital impact in the making of drugs, agrochemicals, flavours, and semi-conducts; they are now and again used both stoichiometrically in examination and business substance responses just as in the job as impetuses. Carbon monoxide responds promptly with any progress metal ions to shape metal carbonyls, themselves a class of organometallics. Metal carbonyls are utilized as catalysts in the petrochemical business.

## **1.1 OVERVIEW OF ORGANOMETALLIC COMPOUNDS AND COMPLEXES**

Most drugs, catalysts, reagents and inhibitors used today are purely organic compounds. Especially after the enormous success of cisplatin in tumour treatment, interest in metal complexes has grown. Synthetic organometallic compounds are generally considered to be toxic or non-compatible with biological systems. Despite this perception, the medicinal properties of organometallic compounds, in particular, organo-transition metal compounds have been probed for a long time and in the last few years, the area has grown considerably (Allardyce and Dyson, 2006).

Catalysts are substances that increase the rate of a reaction but are not consumed in the reaction. Catalysts are widely encountered in nature, industry, and the laboratory (Gupta *et al.*, 2013). Many of the catalysts utilized in the chemical industry and the laboratory are organometallic compounds which are shown in the reactions below:

### **1.1.1 Reaction with Metal and Transmetallation**

The net reaction of an electropositive metal M and a halogen-substituted hydrocarbon is; if one metal atom takes the place of another, it is called transmetallation. Transmetallation is favourable when the displacing metal is higher in the electrochemical series than the displaced metal.

### **1.1.2 Metathesis**

The metathesis of an organometallic compound (MR) and a parallel halide (EX) is a generally utilized engineered course in organometallic science (Gupta *et al.*, 2013). Metathesis response can habitually be anticipated from one or the other electronegativity or corrosive base contemplations. Hydrocarbon bunches will in general cling to the more electronegative

component; the halogen favours the arrangement of ionic mixtures with the more electropositive metal. In short, the alkyl and aryl bunch will in general move from the less to the more electronegative component [X= electronegativity] (Gupta et al., 2013). When the electronegativities are comparable, the right result might be anticipated, with care, by thinking about the mix of the gentler component with natural gathering and harder component with fluoride or chloride. An insoluble item or reactant might change the result, e.g.; HgPhBr ends up being insoluble in THF. Metathesis responses including a similar focal component are regularly alluded to as reallocation responses. Aluminum (Al) is more electropositive than Germanium (Ge), this response happens as it is thermodynamically ideal.

## 1.2 PREPARATIONS OF ORGANOMETALLIC COMPOUNDS


### 1.2.1 Metals with Organic Halides

The planning with metals and natural halides of which its response produces dormant, somewhat polar conditions in which organometallic compounds are regularly solvent. Dampness, oxygen, and carbon dioxide, which respond with it, should be stayed away from. Lithium, magnesium, and zinc are among the metals utilized.



The reactivity request of the halides is  $I > Br > Cl \gg F$ . though magnesium and lithium respond well with chlorides, bromides and iodides. Mercury possibly responds when amalgamated with sodium.

### 1.2.2 Displacement of One Metal by Another

The balance is to such an extent that the R bunch favours the connection of the more electropositive metal.



### 1.2.3 Organometallic Compounds with Metal Halides



The harmony favours the items with R associated with the less electropositive metal so the response will in general shape a less receptive organometallic compound from a more responsive one.

#### 1.2.4 Organometallic Compounds from Acidic Hydrocarbons

Some organometallic compounds are arranged best by the response of a solid base or an alkyl metal subsidiary with an acidic hydrocarbon, like an alkyne.

A particularly significant model is that of 1,3-cyclopentadiene which is acidic because its form base (cyclopentadiene anion) is extraordinarily balanced out by electron delocalization. The anion I framed effectively from the hydrocarbon and methyl lithium.

### 1.3 USES AND APPLICATIONS OF ORGANOMETALLIC COMPOUNDS

- i. It has been utilized as pesticides and fungicides.
- ii. Organometallic intensifies as organotins have for quite some time been utilized as antifouling biocides for boats and fishing nets and rat anti-agents.
- iii. They are utilized in the creation of light-transmitting diode LEDs.
- iv. Utilized in the creation of margarine.
- v. Naturally, there are two uses of organometallic intensifies which are central to human endurance:
  - o Haemoglobin-for moving oxygen in blood
  - o Chlorophyll-for transforming CO<sub>2</sub> into oxygen and natural matter.
- vi. Used in the treatment of cancer.
- vii. Alkene hydrogenation [Wilkinson's catalysis]: Hydrogenation responses are widely used to make business merchandise, hydrogenation is utilized in the food business to make a huge assortment of produced products, similar to spreads and shortenings, from fluid oils. This cycle likewise expands the compound soundness of items and yields semisolid items like margarine. Hydrogenation is additionally utilized in coal preparing. Strong coal is changed over to a fluid through the expansion of hydrogen. Condensing coal makes it accessible to be utilized as fuel. An illustration of an alkenes expansion response is an interaction called hydrogenation. In a hydrogenation response, two hydrogen iotas are added across the twofold obligation of alkenes, bringing about immersed alkenes. Hydrogenation of a

twofold bond is a thermodynamically positive response since it frames a steadier (lower energy) item within the sight of impetus like palladium, nickel and platinum. The Rhodium impetus has been demonstrated to be a reasonable advertiser for the actuation of C-H securities, which has arisen as a difficult and alluring instrument for catalysis





Figure 4: Hydrogenation of Ethene Using Palladium as Catalyst.

- viii. Hydroformylationoxo measure: includes the expansion of carbon monoxide and hydrogen to an alkene to shape an aldehyde containing at least one carbon iotas than the first alkenes. This synergist response is utilized in the petrochemical business, in 1960, Rhodium fills in as exceptionally dynamic impetuses utilized. Since 1970, most hydroformylation depends on impetuses dependent on Rhodium. The resulting research prompted the improvement of water-solvent impetuses that work with the partition of the items from the impetus. The synergist cycle continues through a progression of organometallic intermediates. The aldehydes delivered by hydroformylation are regularly diminished to alcohols that are utilized as solvents, as plasticizers, and in the blend of cleansers.



- ix. Polymerization Alkenes Polymerization Reaction: Polyalkenes, the most widely recognized and valuable class of engineered polymers, are frequently ready by utilization of organometallic impetuses, either in arrangement or upheld in a strong surface. In1950,

the German scientist Karl Ziegler fostered an impetus for ethene polymerization dependent on an impetus shaped by the response of  $TiCl_4$  with  $Al(C_2H_5)_3$ .

Before long, Giulio Natta utilized this kind of impetus for the polymerization of propene to deliver polymers with profoundly standard designs. The private subtleties of the responses of these business synergist measures are not altogether seen, however, there are solid signs from all the more effortlessly considered dissolvable organometallic impetuses that alkenes arrange to a metal place and afterwards embed into a hydrocarbon chain, delivering a more extended chain hydrocarbon connected to the metal community. Reiteration of this cycle prompts amazingly long-chain hydrocarbon polymers, which incorporate a considerable lot of the most natural plastics, like polyethylene and polypropylene. These plastics are utilized in purchaser things going from milk compartments and plastic sacks to fake appendages and vehicle guards.

In catalysis, silver is dynamic as silver oxide in the Wolff revision. Silver nitrate is utilized to isolate out alkenes as the  $\eta^2$ -alkene complex. Silver is additionally present in other carbon-carbon bond skeletal revisions, for example, the quadricyclane to norbornadiene adjustment, the cubane to cuneane reworking and the modification of the cyclobutadiene dimer to cyclo-octatetraene.

- x. Manufacture of acetic acid (Monsanto acetic measure).
- xi. Used in therapeutic medicines.

#### **1.4 Antimalarial Activity of Organometallic Analogs of Known Antimalarial Drugs**

##### **Ferroquine (FQ)**

Ferroquine was found to have an extraordinary result in both CQ-safe and CQ-touchy *P. falciparum* given its capacity not exclusively to repress the hemozoin arrangement, yet additionally to show no insusceptible harmful impacts in rodent models. FQ antimalarial movement came from the association of hydrogen in its parallel side chain which helps keep its mathematical shape set up. Specialists further broke down that the isosterism a piece of FQ had no cooperation with the receptor except for supported the general direction and functionalities of the atom. Furthermore, FQ goes about as a harmful heme crystallization preventer by a social event in the stomach related vacuoles of the parasite, making harm the hemozoin layer, which

brings about the passing of the parasite. Moreover, FQ creates a lot of hydroxyl revolutionaries in the stomach related vacuoles of the parasite, making the parasite kick the bucket. At the point when FQ is in the stomach related vacuole of the malarial parasite, it goes through a reversible one-electron redox response, bringing about the development of salt and hydroxyl extremists. This prompts the development of FQ bunches near the film of the stomach related vacuole, bringing about the arrangement of ROS and lipid peroxidation, prompting the demise of the parasite. FQ capacity to diminish CD4<sup>+</sup> CD25<sup>+</sup> T-cells through fast activity/impact against blood parasites and its capacity to keep up with the multiplication of spleen cells within the sight of various mitogens have driven analysts to recommend FQ as an incredible option in contrast to the treatment of *P. falciparum*.

FQ antimalarial movement has happened to intrigue the scientists and an extension in finding comparative FQ subordinates having antimalarial exercises. A few classes of FQ subsidiaries incorporate hydroxyferroquines, trioxaferroquines, thiosemicarbazone subordinates, ferrocene double forms, chloroquine-connected ferrocenophanes, and 4-N-subbed subordinates.

#### **1.4.1 Hydroxyferroquine**

Subsidiaries. Three hydroxyferroquine subsidiaries were incorporated by Biot et al. to intently impersonate hydroxychloroquine. Although subsidiaries of FQ performed lesser than the first FQ, they were more fruitful in the restraint of in vitro development of *P. falciparum* than CQ. Besides, they showed antiviral impacts in extreme intense respiratory disorder Covid. It has likewise been discovered that ferroquine and subsidiaries have antitumor action. Writing has shown that ferroquine upsets lysosomal work and accordingly restrains autophagy.

#### **1.4.2 Trioxaferroquines**

Trioxaferroquines were accounted for as potential antimalarial specialists against CQ-safe strains, FcB1 and FcM29. Trioxaferroquines were displayed to have high action and extraordinary capacity to clear parasitemia beneath perceptible levels in mice when treated with 10 mg/kg. Subsequently, Trioxaferroquines have been shown as another class of antiparasitic specialists in vivo and in vitro.

### **1.4.3 Thiosemicarbazone Derivatives**

Thiosemicarbazones (TSCs) and FQ analogs show an incredible action against various strains of *P. falciparum*. These analogs amass in food vacuole and showed a preferable parasitic movement over chemical restraint. The utilization of TSCs and FQ showed that there might be more than one method of activity. The first is utilizing falcipain-2 as a sub-atomic objective and the second is acting freely.

### **1.4.4 Ferrocene Dual Conjugates**

These double FQ inferred models showed a superior action in vitro against NF54 and K1 clones of *P. falciparum*. In any case, the double FQ subordinate's movement was lower than that of the parent FQ analogs. This might have been the aftereffect of oxidative action in the stomach related vacuole.

## **2.0 CANCER**

Cancer is a wide term that alludes to an illness that happens when natural anomalies bring about unregulated cell development and division. A few cancer advance quick cell expansion, while others cause cells to isolate and grow all the more leisurely. A few examples like leukemia, produce apparent tumours, while others, for example, breast cancer, don't. The greater part of the body's cells has explicit capacities and fixed life expectancies. While it might seem like something terrible, cell passing is important for a characteristic and helpful marvel called apoptosis. A cell gets directions to bite the dust so the body can supplant it with a fresher cell that capacities are better. Carcinogenic cells come up short on the segments that train them to quit isolating and kick the bucket. Accordingly, they develop in the body, utilizing oxygen and supplements that would as rule support different cells. Carcinogenic cells can shape tumours, hinder the safe framework and cause different changes that keep the body from working routinely. Malignant cells might show up in one region, then, at that point spread using the lymph hubs. These are groups of insusceptible cells situated all through the body.

Specialists arrange cancer by its area in the body or the tissues that it shapes in. For example, sarcomas create in bones or delicate tissues, while carcinomas structure in cells that cover interior or outer surfaces in the body. Basal cell carcinomas create in the skin, while adenocarcinomas can frame in the bosom. At the point when harmful cells spread to different

pieces of the body, the clinical term for this is metastasis. An individual can likewise have more than each sort of malignant growth in turn.

## **2.1 Types of Cancer**

There are over 100 sorts of cancer, beneath are the most well-known sorts:

- i. Bladder
- ii. Lung
- iii. Prostate
- iv. Colon and Rectal
- v. Endometrial
- vi. Kidney
- vii. Leukemia
- viii. Liver
- ix. Melanoma
- x. Non-Hodgkin's Lymphoma
- xi. Pancreatic
- xii. Thyroid
- xiii. Breast

## **2.2 Causes of Cancer**

A portion of the reasons for malignancy incorporates

- i. Smoking
- ii. weighty liquor utilization
- iii. Excess body weight
- iv. actual latency
- v. Malnutrition

## **2.3 Treatments of Cancer**

Specialists for the most part endorse therapies dependent on the sort of disease, its stage at analysis, and the individual's general wellbeing. The following are instances of ways to deal with malignant growth treatment:

- i. Chemotherapy means killing dangerous cells with drugs that target quickly partitioning cells. The medications can likewise assist with contracting tumours, however, the incidental effects can be serious.
- ii. Chemical treatment includes taking medication that changes how certain chemicals function or meddle with the body's capacity to create them. At the point when chemicals assume a critical part, similarly as with prostate and breast cancer, this is a typical methodology.
- iii. Immunotherapy utilizes prescriptions and different therapies to help the insusceptible framework and urge it to battle malignant cells. Two instances of these medicines are designated spot inhibitors and supportive cell move.
- iv. Accuracy medication, or customized medication, is a fresher, creating approach. It includes utilizing hereditary testing to decide the best therapies for an individual's specific show of malignant growth. Specialists still can't seem to show that it can viably treat a wide range of malignancy, nonetheless.
- v. Radiation treatment utilizes high-portion radiation to kill carcinogenic cells. Additionally, a specialist might prescribe utilizing radiation to recoil a tumour before a medical procedure or lessen tumour-related manifestations.
- vi. Undeveloped cell relocation can be particularly advantageous for individuals with blood-related tumours, like leukemia or lymphoma. It includes eliminating cells, like red or white platelets, that chemotherapy or radiation has annihilated. Lab professionals then, at that point reinforce the cells and set them back into the body.
- vii. A medical procedure is frequently a piece of a therapy plan when an individual has a malignant tumour. Likewise, a specialist might eliminate lymph hubs to decrease or forestall the infection's spread.
- viii. Designated treatments perform capacities inside harmful cells to keep them from duplicating. They can likewise support the safe framework. Two instances of these treatments are little particle medications and monoclonal antibodies.

### 3.0 ORGANOMETALLIC COMPOUNDS IN CANCER TREATMENT

An organometallic compound is used in the synthesis of tamoxifen which is taken by those with breast cancer. The activities of the tamoxifen compound are linked to the reversible redox reaction of its iron center. The Grignard reagent is the molecule that is used to add a benzene group to tamoxifen compound, this is because it creates a new bond between two carbons. It was such a big deal in chemistry that François Auguste Victor Grignard won the Nobel prize in chemistry, when carbon is bonded to a more electropositive atom-like magnesium, it possesses a partial negative charge. An organometallic complex is made up of many distinct metals bonded to a carbon atom. Organomagnesium and organolithium complexes are the most prevalent forms of organometallic compounds.

A study from Jaouen and co-workers focusing on prostate cancer therapy used ferrocenyl complexes which possess strong anti-proliferative activity against hormone-independent prostate cancer cells. Ying Li research group also reported novel ferrocenyl derivatives with anti-cancer effects in human lung cancer cells. All of these studies neatly demonstrate the diversity of ferrocenes and their efficacy in various biological profiles as prospective anticancer medicines. Organometallic compounds and complexes utilized in cancer therapy include titanocenes, metallocenes, M-arenes, Ru Arenes, M-carbenes, M-carbonyls, and M-alkyls.

Nanotechnology-derived structures capable of delivering a significant amount of medication to tumour cells in a focused directed procedure have been proposed and developed for application in cancer treatment. Cancer nanotechnology offers a wealth of safety tools for the treatment and diagnosis of cancer, such as targeted devices with unique therapeutic properties that, because of their small sizes can pass biological barriers and deliver multiple therapeutic agents directly to cancer cells and adjacent tissues around the micro-environment.

Nano-vectorization systems have several advantages over traditional chemotherapy, including the ability to hold multiple targeting moieties for multiple target molecules in cancer tissues at the same time, as well as the ability to deliver much larger therapeutic payloads per target while avoiding many biological barriers. Subsequently, titanocene-based organometallic complexes entered clinical trials and their mechanism of action was found to be different to that of the clinically approved drug cisplatin., scientists have focused more on the development of organometallic transition metals antitumor drug candidates since these complexes show high kinetic stability and reactivity due to the presence of a metal-carbon bond and pi-bond arenes.

Similarly attaching a ferrocene moiety in place of the phenyl ring in tamoxifen led to the formation of ferrocifen, which showed improved activity in breast cancer cells and will enter a clinical trial soon. Therefore, the use of organometallic complexes provides much scope for rational drug design with better cellular uptake, minimum unwanted side effects and different modes of action inside tumour cells. These molecules are less toxic and show different modes of action than cisplatin and directly interact with the biological target DNA and undergo ligand hydrolysis.

Besides, organometallic complexes show different modes of action compared to classic metal-based drugs. These complexes show enhanced activity towards cancer cells compared to normal coordination complexes due to the following reasons:

- Direct binding with a target site
- Indirect interaction through non-covalent interactions at the Target
- Activity is mainly due to the metal-arene system, where the metal acts as a carrier *in vivo*
- Production of reactive oxygen species (ROS)
- Organometallic complexes are photoactive and act as a photosensitizer
- Selectivity and efficiency towards multiple targets such as thioredoxin reductase (TrRx), mitochondria, endoplasmic reticulum, DNA, proteins and enzymes
- Controlled ligand substitution and redox reactions and
- Kinetic stability and relative lipophilicity and the presence of a metal in a low oxidation state.

For many centuries, the use of metallodrugs for cancer chemotherapy has been the focus of many scientists. Currently, with the use of modern technology, a better understanding of cancer mechanisms, the strategies for the development of new multifunctional chemotherapeutic drugs have accelerated with less toxicity and different mechanisms of action.

### **3.1 Anticancer Agents**

The creation of metals like cisplatin and carboplatin, which have platinum as a center ion, massively affects current disease chemotherapy. Cisplatin has gotten quite possibly the most broadly used meds in the treatment of illnesses like ovarian and testicular tumours. The restrictions of cisplatin have animated examination in the field of platinum antitumor science by

remembering the decrease for poisonousness of cisplatin (queasiness, ear harm, retching, loss of sensation in hands, and kidney harmfulness), gained drug obstruction saw in specific tumours and failure of the medication against probably the commonest tumours (e.g., colon and breast). Because of its specific compound construction, cisplatin offers little opportunities for development in tumour explicitness and subsequently diminishing incidental effects. The other option edifices have no less than one immediate, covalent metal-carbon bond, having underlying assortment, different stereochemistry, give power over major active properties, dynamically steady, normally uncharged, and having low oxidation condition of metal ion, so they can be utilized as an optimal contender for anticancer competitors. A few models are metallocenes, organometallic ruthenium half-sandwich buildings, organometallic osmium edifices half-sandwich buildings, organometallic iridium and rhodium buildings, rhenium organometallics, ruthenium, osmium, iridium, and platinum organometallics as frameworks for protein kinase inhibitors, metal NHC edifices, and metal carbonyl edifices.

### **3.2 Antibacterial Agents**

The greatest test in the antibacterial market is the issues identified with the medication safe microbes. The cure is presently to look for new mixtures with new methods of activity to beat the safe strains. This should be possible by either the natural derivatization of old medications or new organometallic drugs, for instance, new tamoxifen, platensimycin, and so on.

### **3.3 Anti-infectant Agents**

Progress metal as silver is being utilized as the antimicrobial specialist because of its low harmfulness when contrasted with different metals. For instance: silver sulfazine, which is utilized to get consumes forestall bacterial diseases. Silver nitrate is given to newborn children to forestall the improvement of ophthalmia neonatorum. Chlorhexidine-silver sulfadiazine is an enemy of infective metal complex against catheter diseases. Organometallic edifices of Pt, Rh, Ir, Pd, and Os metal with dynamic natural particles have been accounted for to show trypanocidal movement. Metal edifices of Pt (II) and Ru (II) with o-vanillin-(4-methyl thiosemicarbazone), metal buildings of Ga (III), Al (III), and Fe are among the different medications.

### **3.4 Anti-inflammatory Agents**

These buildings are likewise utilized as calming and against ligament specialists. A few injectable progress metal buildings as sodium aurothiomalate, aurothioglucose, sodium aurothiopropanol, and gold and silver nanoparticles formed with heparin subsidiary have antiangiogenesis properties. Gold has been utilized for the treatment of fringe psoriatic arthropathy. As a result of oxygen digestion, a superoxide anion can trigger oxidative injury to tissues. This action is related to ripper combination and provocative sicknesses just as neurological issues like Parkinson's illness and Alzheimer's infection. In any case, overabundance utilization of these buildings in joint inflammation causes agony and fever. Nitrosyl (NO) is a brilliant ligand for progress metal particles and these metal nitrosyls having helpful qualities. Sodium nitroprusside is utilized to treat cardiovascular sicknesses by delivering NO with restricted utilization because of the harmfulness of CN<sup>-</sup>. Ruthenium poly amino-carboxylate buildings are likewise proficient NO foragers.

### **3.5 Anti-diabetic agents**

Diabetes is the most enduring illness among individuals. This is the infection wherein the body doesn't deliver insulin chemical, which is utilized for retention of glucose in cells. The control of glucose levels is finished by vanadium buildings with natural ligands which are less poisonous and have further developed dissolvability and lipophilicity. These edifices show association in the initiation of conspicuous key segments of insulin flagging pathways. Chromium supplementation likewise improves glycemia among patients with diabetes. Additionally, higher zinc consumption likewise brings down the danger of type 2 diabetes in ladies.

### **3.6 Neurological Drugs**

Change metal buildings are additionally used to treat neurological sicknesses. Huntington's chorea, tardive dyskinesia, convulsive torticollis, Tourette's condition, L-dopa instigated hyperkinesia, Parkinsonism, natural cerebrum problems, drug prompted capricious issues, headache and group migraine, occasional hyper-lethargy, epilepsy, Meniere's illness, and intermittent hypokalemic loss of motion are among the conditions for which lithium is recommended. Lithium hinders the rummaging pathways for catching inositol in the resynthesis of poly phosphoinositides in the cerebrum. Zinc is additionally utilized as a transmitter in neuronal flagging pathways.

### 3.7 Delivery Probes and Diagnostic Tools

Organometallic edifices have novel properties as redox movement, Lewis acidity, electrophilicity, valency, calculation attractive spectroscopic, and radiochemical properties which can be utilized to gauge cell capacities. Gold nanorods have been utilized for photoacoustic sub-atomic imaging with concurrent different focusing as they are less responsive and less poisonous. The nanoparticles infused in the tumour cells build their capacity to ingest radiation of explicit frequency. The property is utilized in lymphotropic nanoparticle upgraded attractive reverberation imaging of prostate disease. As iron oxide has superparamagnetic properties so it can go about as a regrettable differentiation specialist in attractive reverberation imaging (MRI) which is utilized to identify the affectability of excited tissues.

Change metals display diverse oxidation states and can associate with a few adversely charged particles. Because of their essential job in restorative science, we have included both the large scale and the nano-organometallic buildings and their underlying and photograph actual conduct exhaustively.

Metallocene compounds have two  $\pi$ -reinforced cyclopentadienyl (Cpa) ligands on a metal particle. These mixtures are likewise called "sandwich buildings" because of their even nature. Other metal edifices with cyclic  $\pi$ -edges are additionally named metallocenes. Mixtures with just one  $\pi$ -edge are named "half sandwich metallocenes". The bis-cyclopentadienyl edifices are partitioned into two classifications: (a) "old style" with equal Cp rings and (b) "bowed" metallocenes, which have different ligands attached to the metal notwithstanding the Cp rings. Ferrocene was the first organometallic compound with antiproliferative properties, so the therapeutic properties of the complex were researched. Ferrocene is a nontoxic compound and can be infused, breathed in, or taken orally. It can't cause significant medical issues. Another Ferrocene-containing compound chloroquine (subordinate) is utilized as an antimalarial drug. Ferroquine has an action like chloroquine on the jungle fever parasite *P. falciparum*. The p-methoxy-benzyl subbed titanocene shows generally excellent action against renal cell malignant growth and pleura mesothelioma cell lines. Ruthenium buildings have low poisonousness and it has a similar system (ligand trade energy) to those of platinum(II) antitumor medications. A class of ruthenium(II)- arene edifices that are feebly cytotoxic in vitro,  $\text{Ru}(\eta^6\text{-p-arene})\text{Cl}_2(1,3,5\text{-triaz-7-phosphaadamantane})$  named as RAPTA, communicate unequivocally with proteins, with

the capacity to segregate restricting to various proteins, however, show a generally low penchant to tie DNA, which is viewed as the fundamental objective of many metal-based medications. Dyson et al. as of late depicted the arrangement of a progression of RAPTA-type buildings with fluoro-subbed  $\eta^6$ -arene ligands. The dynamic Pt-drug is by all accounts cis-facilitated by bidentate amine ligands or two amines (no less than one – NH bunch on the amine) and two leaving bunches with a middle of the road restricting strength (e.g., Cl<sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, citrate or oxalate) to platinum. Non-platinum metals might have some unique substance conduct (oxidation state, redox potential, coordination calculation, extra coordination locales, restricting inclinations to biomolecules as the HSAB (hard and delicate (Lewis) acids and bases), the pace of hydrolysis or energy of ligand trade responses and the capacity to supplant fundamental metals. Because of these distinctions, the non-platinum metal-based mixtures might have various components of activity, bio-conveyance, and organic movement.

NESTEC 2021

## CONCLUSION

Conclusively, it was established that organometallic compounds and complexes have therapeutic potential towards curing cancer, diabetes, HIV/AIDS, malaria and Chagas disease that is clinical while industrially many chemical organic compounds such as alcohols, acetic acids, polyethylene and polypropylene are produced from these complexes. Industrial and clinical uses of organometallic complexes and some prominent studies on new therapeutic complexes were mentioned. Developments in explorations of organometallic compounds, in various therapeutic areas, are continued to be an active and productive area of research. Increasingly powerful tools, notably spectroscopic techniques like time-resolved infrared are used for identifying and structurally characterizing the solid complex. Optical spectroscopic and potentiometric methods are used for monitoring intermediates and species in solution. Besides these preclinical and clinical studies significantly enhanced our knowledge and understanding of the structure and mechanistic aspects of the therapeutic of organometallic complexes. The dynamic properties of organometallic compounds and complexes are often probed with variable temperature, nuclear magnetic resonance (NMR) and chemical kinetics hence help in many vital applications both industrially and clinically.

We can identify practically any 'traditional' organometallic chemical class to have been tested for biological activity, thanks to the expanding number of researchers entering the field of bio organometallic chemistry.

The biggest change in the drug development in the anti-cancer industry has been the move from cytotoxic to molecularly targeted agents, as nano-medicine has brought about new horizons to cancer therapy. Promising organometallic drug candidates can now be selectively delivered to tumour cells with minimum side effects.

Note, not all compounds containing carbon and metal atoms are organometallic compounds.

## REFERENCES

- Benjamin, inc. Menlo Park, CA. ISBN 0-8053-8329-8.
- Biot, C.G.; Dive, D. In: Medical Organometallic Chemistry; Jaouen, G.; Metzler-Nolte, N., Eds., Springer Berlin Heidelberg: Berlin, Heidelberg, 2010, Vol. 32. Pp. 155-193.
- Comprehensive organometallic chemistry II by F. Gordon Stone; Geoffrey Wilkinson; Edward W. Abel Gupta, B. D., Elias, Anil J. (2013). "Basic Organometallic Chemistry: Concepts, Syntheses and Applications". Universities Press, CRC Press. ISBN 978-81-7371-8748.
- Dive, D; Biot, C. Ferrocene conjugates of chloroquine and other antimalarials: the development of ferroquine, a new antimalarial ChemMedChem 2008, 3, 383-391.
- Dr O.A Azeez, chemical engineering department, Futminna, Niger state.
- Gupta, B. D., Anil, S., and Elias, J. (2013). Basic Organometallic Chemistry: Concepts, Syntheses and Applications Publisher Universities Press, CRC Press ISBN=978-81-7371-8748.
- <https://www.britannica.com>
- Jaouen, G.; Dyson, P.J. Medicinal Organometallic Chemistry. In Comprehensive Organometallic Chemistry III. From Fundamentals to Applications; Crabtree, R. H.; Mingos, D. M. P., Eds; Elsevier: Oxford, U.K., 2007; Vol. 12, pp 445.
- Jaouen, G., Ed. Bioorganometallics: Biomolecules, Labeling, Medicine; Wiley-VCH: Weinheim, Germany, 2006.
- John D. Robert and Marjorie C. Caserio (1977) Basic Principle of organic chemistry, second edition W.A.
- Kopf-Maier, P; Hesse, B, Voigtlander, R.; Kopf, H. Tumor inhibition by metallocenes: antitumor activity of titanocenedihalides ( $C_5H_5$ )<sub>2</sub>TiX<sub>2</sub> (X=F, Cl, Br, I, NCS) and their application in buffered solutions as a method for suppressing drug-induced side effects J.Cancer Res. Clin. Oncol. 1980, 97, 31-39.
- Osakada, Kohtaro (2003). Fundamentals of Molecular Catalysis. Amsterdam: Elsevier. ISBN 0444509216.
- Periana, R.A., Bhalla, G., Tenn, W.J., III, Young, K.J., H., Liu, X.Y., Mironov, O., Jones, C., Ziatdinov, V.R. (2004). Perspectives on some challenges and approaches for developing the next generation of selective, low temperature, oxidation catalysts for alkane hydroxylation based on the C-H activation reaction. Journal of Molecular Catalysis A: Chemical, 220 (1) 7-25.